


INFORMATION LEAFLET FOR PARENTS / CARERS OF A CHILD

Volumatic Education Action Plan

Older than 6 years old



Volumatic (Spacer Device)

Spacers act as a holding chamber for inhaled medication from a metered dose inhaler, allowing time for medication to be inhaled.

1. Fit the two halves of the volumatic together.
2. Remove the cap of the inhaler.
3. Shake the inhaler well.
4. Fit the inhaler firmly into the adaptor end of the volumatic.
5. Place the mouthpiece in the mouth between lips and teeth.
6. Seal the lips firmly around it. Avoid putting lips over the side holes of the mouthpiece.
7. **Breathe out** into the device as far as is comfortable.
8. Press the inhaler once to release a dose
9. Breathe in and out slowly and deeply five times. The valve should click with each breath

For every dose the inhaler should be removed from the volumatic and steps 3 -9 repeated.

Cleaning (needs to be replaced 9-12 monthly)

- Clean twice weekly. Pull the halves gently apart. Do not take the valve apart to reduce static.
- Soak in warm water and mild detergent for a few minutes. Use a soft toothbrush to remove any film on the valve or inside the spacer device.
- Do not rinse or rub the spacer device leave to dry at room temperature rinsing or rubbing will create static.
- Wipe the mouthpiece to remove detergent before use.
- Change volumatic if valve gets damage or fails to open.



If Inhaler hasn't been used in a week or more it may need to be primed, do so by releasing one puff into the air.

Only release one puff of the inhaler into the volumatic at one time.

Volumatics are single person use and should not be shared.

For further information and demonstrations on inhaler devices: www.asthmasociety.ie

General Paediatric Team December 2020

Developed by General Paediatrics – Clinical Nurse Specialists – Date issued 03.12.20

Disclaimer & Copyright ©2019, Children's Health Ireland at Crumlin, Dublin 12. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior written permission of the copyright holder. Every effort has been made to ensure that the information provided is accurate and in accord with standards accepted at the time of printing.



My Asthma Medicine

- My Controller inhaler is.....
- Colour.....
- I take puff/s of my preventer / controller in the morning and.....puffs in the evening.
- My oral controller is.....
- I take my Controller medicine all the time, even when well.
(Don't forget to rinse mouth after using controller)
- *My Controller benefits my lungs by reducing inflammation, swelling and mucus.*
- My Reliever Medication is.....
- Colour.....

ALWAYS USE SPACER DEVICE WHEN USING INHALERS

My Asthma Action Plan		
Orange Zone	Green Zone	Red Zone
Feeling Unwell	Feeling Well	Feeling Worried
I cough or wheeze and it is hard to breathe	Feeling well, no symptoms	My reliever is not helping
I am waking at night because of my asthma	I have no cough	I am finding it hard to breathe and / or talk or feed
I cough or wheeze when I play	I play just like other children	I am breathing hard and fast
I need my reliever inhaler to control my asthma more than 2 times a week	I use my reliever less than 2 times a week	I am sucking in around my ribs / throat
		I am looking pale or blue
I continue to take my controller inhaler every day	I take my Controller Inhaler EVERY day.	My reliever is having little or no effect
If my symptoms are not improving or getting worse, contact GP or Emergency Department	2 puffs via spacer Device	I need to take one puff of my reliever inhaler every 30-60 seconds – up to a total of 10 puffs.
		Take slow and steady breaths
		Stay Calm and seek Urgent medical advice/GP or Emergency Department or call an ambulance

Developed by General Paediatrics – Clinical Nurse Specialists – Date issued 03.12.20