

NURSING CARE PLAN No 27b
Enteral Feeding
All careplans must be used in conjunction with careplan 1

Full Name:

Address: **Addressograph**

HCR:.....

Care Plan No 27b Problem	Enteral Feeding Goals	Issue Date: April 2020 Review Date: April 2023
---	--	---

<p>..... requires Enteral feeding via</p> <p><input type="checkbox"/> Nasogastric</p> <p><input type="checkbox"/> Naso Jejunal</p> <p><input type="checkbox"/> Mic-Key Tube</p> <p><input type="checkbox"/> PEG Tube</p> <p><input type="checkbox"/> Mallecot / Foley gastrostomy tube</p> <p><input type="checkbox"/> Transgastric Jejunal</p> <p>Date tube inserted:</p>	<ul style="list-style-type: none"> Enteral feeding will be managed appropriately Discharge Safely
--	---

Use in conjunction with OLCHC Guidelines on the Management of Enteral Feeding 2nd edition Feeding via Peg, Mic-Key

NB. Loop all enteral feeding tubes when not in use to prevent tubes catching in equipment if not connected to feeding pumps. Special care when moving and handling infants

(For example, transferring to a buggy, placement in parents' arms or weighing)

No	NURSING INTERVENTION	Commencement, Date, Signature, Time, Grade	Discontinued, Date, time, Signature, Grade
----	----------------------	--	--

1a	Naso-Gastric (NG) tube		
<ul style="list-style-type: none"> Use in conjunction with NPC guidelines on Nasogastric tube placement Check pH prior to each feed or medication administration -range between 0-5.0 If pH not within normal limits, please refer to NG Guidelines & to the Medical / surgical Team Flush tube with 10ml syringe with 10ml of sterile water pre and post feed/medication administration (Adjust flushing volume as per child i.e. fluid restricted) Change tube as per manufactures instructions 			
Date Tube Inserted		Date Tube Inserted	
Size		Size	
Length		Length	

1b	Naso-jejunal (NJ) Tube		
<ul style="list-style-type: none"> Position of NJT confirmed by X-Ray PFA (Plain Film Abdomen) Nurses can only pass NJT to stomach it needs to be advanced further by the Medical/Surgical Team Measure NJT tube from nare to tube end following insertion and confirmed in place At the commencement of each shift measure tube from nare to tube end prior to any feeding /medication administration Ensure measurements compare accurately with measurements taken on previous shifts Length _____ cms Date _____ pH is not an indicator of NJT tube position as the jejunal aspirate should be alkaline. An aspirate may be requested by the team to ensure the tube is not in the stomach (indicated by a pH value between 0-5). ANTT level 2 must be used prior to the connection of NJ T. The Jejunum does not contain the same protective barrier as the gastric acid and therefore infection is more likely. 			

NURSING CARE PLAN No 27b
Enteral Feeding
All careplans must be used in conjunction with careplan 1

Full Name:

Address: **Addressograph**

HCR.....

<ul style="list-style-type: none"> Avoid granular medications such as Omeprazole and Lansoprazole as such medications have the potential to block the tube if such medication is required, please discuss the need for liquid suspension. 			
1c	Mic-Key Tube	Commencement , Date, Signature, Time, Grade	Discontinued, Date, time, Signature, Grade
<ul style="list-style-type: none"> Check pH prior to the first feed - Range should be between 0 – 5. Rotate Mic–Key 360 degrees, 72 hours post insertion and continue this rotation daily If displacement of the tube occurs check the pH and refer to the Medical/Surgical Team Change water in balloon every 7 days, remove the 5mls of water with a 10ml syringe and replace with 5mls of water in a 5ml syringe Supplies out of hours: If supplies needed outside of hours, these are available in Materials Management Storeroom (Bleep 8700) contact security for access to same 			
1d	PEG Tube		
<ul style="list-style-type: none"> Check pH prior to the first feed - Range should be between 0 – 5. Rotate the PEG tube 360 degrees 72hrs post insertion Rotate tube daily to ensure tract formation and prevent Buried Bumper Syndrome If displacement of tube occurs check pH and refer to the Medical/Surgical Team <p>Additional information:</p>			
1e	Mallecot / Foley Gastrostomy		
<ul style="list-style-type: none"> DO NOT ROTATE TUBE (<i>Tube may be held in place by a suture as well as a water balloon</i>) Gastric pH MUST be checked before each use. Range should be between 0 – 5. Document same in nursing notes. <p>Additional information:</p>			

NURSING CARE PLAN No 27b
Enteral Feeding
All careplans must be used in conjunction with careplan 1

Full Name:

Address: **Addressograph**

HCR.....

1f		Commencement , Date, Signature, Time, Grade	Discontinued, Date, time, Signature, Grade
	<p>DO NOT ROTATE TUBE</p> <p style="text-align: center;">pH must be checked prior to first feed</p> <p style="text-align: center;">pH reading from the Gastric port < or = 5.0</p> <p style="text-align: center;">pH reading from the Jejunal port > or = 6.0</p> <ul style="list-style-type: none"> • Continuous feeds only through the jejunal port to prevent dumping syndrome • Use ANTT Level 2 when setting up feed or attending to the tube • Use 10ml syringe only when flushing pre and post feeds (maximum volume of flush 10ml and minimum volume 5mls) • If jejunal port is not in use daily, it must be flushed twice daily with 10ml of sterile water to avoid blockages. • If jejunal port is used for feeds gastric outlet must be flushed daily with 10ml sterile water in a 10ml syringe. • Avoid granular medications such as Omeprazole and Lansoprazole as they have the potential to block the tube. If such a medication is required discuss the need for liquid suspension. Clarithromycin should be avoided as such medication can block tube completely. 		
2	Discharge Planning		
	<ul style="list-style-type: none"> • Provide child/parent/guardian with written and verbal information (<i>Advice for the parents/guardians caring for an infant/child with a Cor-Flo Percutaneous Endoscopic Gastrostomy PEG Tube/Mic-Key Button</i>) • Provide education to child/parents/guardians and ensure Parent Teaching Checklist/Competency Assessment is completed prior to discharge. <p>Mic-Key Tube only</p> <ul style="list-style-type: none"> • Advise child/parents/guardian to make appointment for three months with Surgical Team to change Mic-Key button • Ensure child/parents/guardians have a spare Mic-Key on discharge which is the same length and size as tube in-situ. • Ensure prescription available to parents/guardian for supplies • Ensure contact is made with the Public Health Nurse for supplies and support post discharge <p>Additional Information</p> <p>.....</p> <p>.....</p>		

Created by Nursing Department
Issue Date: April 2020 / Review Date: April 2023

Copyright and Disclaimer @2020. Children's Health Ireland at Crumlin, Dublin 12. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior written permission of the copyright holder. Every effort has been made to ensure that the information provided is accurate and in accord with standards accepted at the time of printing.