

FLACC Behavioural Pain Assessment Scale

CATEGORIES	0	1	2
Face	No particular expression or smile	Occasional grimace or frown; withdrawn, disinterested	Frequent to constant frown, clenched jaw, quivering chin
Legs	Normal position or relaxed	Uneasy, restless, tense	Kicking or legs drawn up
Activity	Lying quietly, normal position, moves easily	Squirming, shifting back and forth, tense	Arched, rigid, or jerking
Cry	No cry (awake or asleep)	Moans or whimpers, occasional complaint	Crying steadily, screams or sobs; frequent complaints
Consolability	Content, relaxed	Reassured by occasional touching, hugging, or being talked to; distractible	Difficult to console or comfort

How to score
In patients who are awake: observe for 1 to 5 minutes or longer. Observe legs and body uncovered. Reposition patient or observe activity. Assess body for tenseness and tone. Initiate consoling interventions if needed.
In patients who are asleep: observe for 5 minutes or longer. Observe body and legs uncovered. If possible, reposition the patient. Touch the body and assess for tenseness and tone.

- Face**
- > Score 0 if the patient has a relaxed face, makes eye contact, shows interest in surroundings.
 - > Score 1 if the patient has a worried facial expression, with eyebrows lowered, eyes partially closed, cheeks raised, mouth pursed.
 - > Score 2 if the patient has deep furrows in the forehead, closed eyes, an open mouth, deep lines around nose and lips.
- Legs**
- > Score 0 if the muscle tone and motion in the limbs are normal.
 - > Score 1 if patient has increased tone, rigidity, or tension; if there is intermittent flexion or extension of the limbs.
 - > Score 2 if patient has hypertonicity, the legs are pulled tight, there is exaggerated flexion or extension of the limbs, tremors.
- Activity**
- > Score 0 if the patient moves easily and freely, normal activity or restrictions.
 - > Score 1 if the patient shifts positions, appears hesitant to move, demonstrates guarding, a tense torso, pressure on a body part.
 - > Score 2 if the patient is in a fixed position, rocking; demonstrates side-to-side head movement or rubbing of a body part.
- Cry**
- > Score 0 if the patient has no cry or moan, awake or asleep.
 - > Score 1 if the patient has occasional moans, cries, whimpers, sighs.
 - > Score 2 if the patient has frequent or continuous moans, cries, grunts.
- Consolability**
- > Score 0 if the patient is calm and does not require consoling.
 - > Score 1 if the patient responds to comfort by touching or talking in 30 seconds to 1 minute.
 - > Score 2 if the patient requires constant comforting or is inconsolable.

Each category is scored on the 0–2 scale, which results in a total score of 0–10.

0: Relaxed and comfortable
1–3: Mild discomfort
4 –6: Moderate pain
7–10: Severe discomfort or pain or both

From Merkel, S. I., Voepel-Lewis, T., Shayevitz, J. R., & Malviya, S. (1997). The FLACC: A behavioral scale for scoring postoperative pain in young children. *Pediatric Nursing*, 23(3), 293–297. The FLACC scale was developed by Sandra Merkel, MS, RN, Terri Voepel-Lewis, MS, RN, and Shobha Malviya, MD, at C. S. Mott Children's Hospital, University of Michigan Health System, Ann Arbor, MI. Used with permission